

Southern Region Clinical Audit Results
2018 – 2019

Contents

Introduction.....	3
Purpose	4
Objectives.....	4
Methodology.....	4
Medical Audit Team:	4
Data Analysis Plan for the Clinical Audit	5
Diabetes and Hypertension:	5
.....	5
Diabetes Results SR.....	6
Outcome Criteria	19
Hypertension Audit Results 2018-2019:	22
Hypertension: By age range:.....	41
General Observations Applicable to all Clinics:	42
Observations per Clinic:.....	43
Dangriga and Satellites (Pomona, Hopkins) Clinical Audit:	43
Diabetes:.....	43
Hypertension:.....	43
Overall:	43
Independence PC and Satellites Clinical Audit:.....	45
Diabetes:.....	45
Hypertension:.....	45
Overall:	45
Punta Gorda and Satellites Clinical Audit:	46
Diabetes:.....	46
Hypertension:.....	46
Overall:	46
San Antonio and Satellites Clinical Audit:	47
Diabetes:.....	47
Hypertension:.....	47
Overall:	47
Bibliography.....	48

Introduction¹:

The National Health Insurance Scheme conducted medical audits at these Primary Care Provider (PCP): **Dangriga, Independence, Punta Gorda and San Antonio**. These PCPs utilize standardized clinical protocols on the management of chronic-non-communicable diseases. The audit tools were modified to capture the key indicators in a more concise manner, facilitating the assessment of overall compliance. The auditors consisted of a team of medical doctors including internists and National Health Insurance representatives. The auditors evaluated clinicians in relation to compliance with clinical protocols implemented by the Ministry of Health for the management of Hypertension and Diabetes Mellitus. Protocols are a common point of reference for prospective and retrospective audits of clinicians' practices: the tests, treatments, treatment goals provide ready process measures for rating compliance with best care practices. Healthcare systems that provide services and government bodies that pay for them have found clinical protocols to be effective in improving efficiency by standardizing care and optimizing value for money.

Non-communicable diseases have become more prevalent in developing countries where the burden of disease is double than that of infectious diseases. If the present trend is maintained, the health systems in low-and middle-income countries will be unable to support the burden of disease. Prominent causes for heart disease, diabetes, cancer and pulmonary diseases can be prevented but urgent preventive actions are needed, and efficient strategies should deal seriously with risk factors like smoking, alcohol, physical inactivity and western diet.

As the Belizean population ages, chronic disease will have an increasing effect on our health and wellbeing. Our health system must therefore be able to respond in an appropriate and cost-effective way to this challenge. Failure to prevent, detect and treat chronic disease at an optimal stage in its course impacts on affected individuals and their families in terms of pain and suffering, and in productivity losses and high health care costs at the community level.

The Central American Diabetes Initiative (CAMDI): Diabetes and Hypertension Risk Factor Survey conducted in Belize in 2006 showed that the overall prevalence of diabetes mellitus was found to be 13.1%. In Belize, diabetes has been reported as the second leading contributor to mortality for 2003 and 2004 and the leading contributor to mortality in 2005 and 2006 and remains the leading cause of death in 2008.

Reports from the Ministry of Health indicate that in 2008, Hypertension was recorded as being the eighth leading cause of death. Hypertension causes organ damage which contributes to ischemic heart disease, the second leading cause of death and cerebro-vascular disease, the sixth leading cause of death in 2008. The study of Diabetes, Hypertension and Non-Communicable Disease Risk Factors published in 2008 indicate that the national prevalence of hypertension among persons twenty years and older was 28.7%.

The results of the 2018-2019 audit also the National Health Insurance with essential data for awarding financial incentives. This incentive is dependent on the achievements of the Key Performance Indicators (KPIs) which are included in the contractual agreement with PCPs. In addition, the analysis of results identifies areas for improvement in the management of Hypertension and Diabetes Mellitus at the Primary Care Level.

Purpose:

- The Medical Audit 2018-2019 was conducted to determine the level to which PCPs provide healthcare services in compliance with the clinical protocols for the management of diabetes and hypertension.

Objectives:

- Conduct clinical audits at the PCPs that provide healthcare services under National Health Insurance.
- Identify factors that create barriers to protocol adherence.
- Propose recommendations based on the findings of audit aimed at improving the quality of NHI services.

Methodology:¹

Sample Selection:

Based on the population size of the registered members within the corresponding PCPs and those identified as diagnosed with either Diabetes or Hypertension, a random sample of 19 records were selected and the Lot Quality Sampling Methodology applied for the analysis.

Medical Audit Team:

The Medical Audit team comprised of a team of doctors:

Compliance with Diabetes and Hypertension Protocols:

Dr. I. Depaz - Internist

Dr. A. Hotchandani – General Practitioner; Consultant

Dr. J. Perez – General Practitioner (QAM NHI)

NHI Personnel:

Dr. N. Castillo – General Manager

Dr. Johanne Perez – Quality Assurance Manager

Ms. Ruth Jaramillo- Health Services Manager

Ms. Cristina Ake- Data Analyst

Mr. J. Colli – I.T. Analyst

The team of doctors participated in the review of the audit tools and in a training on the application of these prior to the audit exercise. The NHI team assisted in the coordination of the activity, development of the audit tool and study design and data analysis.

Data Analysis Plan for the Clinical Audit¹:

Diabetes and Hypertension:

The emphasis of the clinical audit exercise was to ensure that critical data necessary to effectively assess compliance with protocols is not only being collected but also that the cases are being managed in accordance to the established protocols for the management of Diabetes and Hypertension.

To determine compliance with the protocol each medical record was assessed in accordance to the audit tool which identifies the following:

1. The **“must do”** criteria or the core variables that should be done. These variables verify that the client has been diagnosed appropriately and that the overall management takes into account, relevant follow up tests and potential target organ damage monitoring. It offers a more holistic approach to the case management of the patient.
2. The **“should do”** criteria seek further input on the supportive management of the client to include nutritional education; renal failure monitoring, importance of adherence and relevance of summary report for continuous clinical management.

The final percentage score was then calculated per record. The minimum passing acceptable score for each record was **90%** or higher.

A percentage was then calculated based on the number of records that met the minimum standard over the number of records assessed. A clinic met the protocol if the percentage of records that met the minimum standard which was also **90%** or higher.

Diabetes Results SR

2018	DAN	IND	PG	SA
Scores (target 90%)	100%	100%	100%	100%

Our clinics continue to perform very well in the clinical audit. All clinics passed on their number of records assessed. This is the consistent pattern that we are seeing. Overall the PCP team have embraced a minimum standard of care that is reflected in their documentation processes.

As we can see below, the documentation that is occurring at the level of the PCPs and their satellites is very good. This reflects the dedication of our teams to give our patients the quality of care that is expected at our facilities:

	DAN	Hopkins	Pomona	Score
# of records assessed	19	5	5	29
# of records that met the target (90%)	19	5	5	29
Total score	100%	100%	100%	100%

	IND	Bella Vista	Placencia	San Juan	Score
# of records assessed	19	5	5	5	34
# of records that met the target (90%)	19	5	5	5	34
Total score	100%	100%	100%	100%	100%

	PG	Big Falls	Santa Ana	Score
# of records assessed	19	5	5	29
# of records that met the target (90%)	19	5	5	29
Total score	100%	100%	100%	100%

	SA	Pueblo Viejo	Santa Teresa	SPColumbia	Score
# of records assessed	19	5	5	5	34
# of records that met the target (90%)	19	5	5	5	34
Total score	100%	100%	100%	100%	100%

Demographics	DAN	IND	PG	SA
Gender				
Male	12	10	4	9
	41%	29%	14%	27%
Female	17	24	25	25
	59%	71%	86%	74%

Females predominantly attend to our clinics (AVG. 72% of the total) in relation to males (AVG. 28%). This is the same ratio as we found in the 2017 audit (AVG. 71% vs 29%). Punta Gorda in particular had a higher percentage of women patients.

Demographics	DAN	IND	PG	SA
Age Range				
31-45	1	6	7	8
	3%	18%	24%	24%
46-55	4	14	5	16
	14%	41%	17%	47%
>55	24	14	15	10
	83%	41%	52%	29%

The average of the total population 31-45 years old has decreased to 17% from 25% (from the 2017-2018 audit). The average population in the 46-55-year-old population increased slightly to 30%, from 28%. And the total average number in the > 55-year-old population increased from 45% to 51%. Dangriga (83%) and SA (53%) had the highest percentage of > 55-year-old patients.

Demographics	DAN	IND	PG	SA
Ethnicity				
Maya	1	5	14	30
	3%	15%	48%	88%
Mestizo	10	17	5	0
	35%	50%	17%	0%
Garifuna	9	4	5	0
	31%	12%	17%	0%
Creole	7	4	1	0
	24%	12%	3%	0%

The predominant ethnic group at all the clinics is Maya (Averaging 39% of the population), followed by Mestizo (AVG. of 25%). San Antonio (88%) and Punta Gorda (48%) had their highest populations being Maya. Garinagu patients are found predominantly in Dangriga and Independence.

Demographics	DAN	IND	PG	SA
Cases of Diabetes				
Newly Diagnose	0	0	0	1
	0%	0%	0%	3%
Established Case	29	34	29	33
	100%	100%	100%	97%

The vast majority of cases currently being seen are established cases.

Demographics	DAN	IND	PG	SA
Classification				
Diabetes Type 1	0	1	0	0
	0%	3%	0%	0%
Diabetes Type 2	29	33	29	34
	100%	97%	100%	100%

Our clinics are encountering type 2 diabetes predominantly. This is the same pattern as in last year's audit.

Diabetes “Must Do” Processes:

As per the tables below:

Our patients are being diagnosed appropriately by all our clinics. We commend the clinics in carrying this out successfully. Clinics are also requesting blood markers of diabetes control, for example, they order the HbA1c minimally once yearly to all their patients. This practice and the search for medication side-effects such as hypoglycemia are engendered in the clinical practice of the teams.

Our patients are being diagnosed appropriately by our clinics. We commend the clinics in carrying this out successfully. Clinics are also requesting blood markers of diabetes control, for example, they order the HbA1c minimally once yearly to all their patients. This practice and the search for medication side-effects such as hypoglycemia are engendered in the clinical practice of the teams.

"Must Do" Process Criteria	DAN	Hopkins	Pomona
The Diagnosis of Diabetes is correct	2.0	2.0	2.0
HbA1c has been checked at least annually	2.0	2.0	2.0
At least annually there has been an assessment of symptoms including hypoglycaemic attack.	2.0	2.0	2.0

"Must Do" Process Criteria	IND	Bella Vista	Placencia	San Juan
The Diagnosis of Diabetes is correct	2.0	2.0	2.0	2.0
HbA1c has been checked at least annually	2.0	2.0	2.0	2.0
At least annually there has been an assessment of symptoms including hypoglycaemic attack.	2.0	2.0	2.0	2.0

"Must Do" Process Criteria	PG	Big Falls	Santa Ana
The Diagnosis of Diabetes is correct	2.0	2.0	2.0
HbA1c has been checked at least annually	2.0	2.0	2.0
At least annually there has been an assessment of symptoms including hypoglycaemic attack.	2.0	2.0	2.0

"Must Do" Process Criteria	SA	Pueblo Viejo	Santa Teresa	SPColumbia
The Diagnosis of Diabetes is correct	2.0	2.0	2.0	2.0
HbA1c has been checked at least annually	2.0	2.0	2.0	2.0
At least annually there has been an assessment of symptoms including hypoglycaemic attack.	2.0	2.0	2.0	2.0

As per the tables below:

Diabetics feet are being well monitored yearly by all the clinics. This process is recorded in all files of diabetics at all clinics. This is a process that is now ingrained in the culture of the clinics.

Microalbuminuria is an important clinical finding because it is not only associated with an increased risk of progression to overt proteinuria (macroalbuminuria) and renal failure but also cardiovascular events. In patients who progress to overt nephropathy, microalbuminuria usually precedes macroalbuminuria by an interval of 5 to 10 years.² Our clinics are requesting microalbumin in urine as a part of their holistic patient diagnosis. Of note, we have re-introduced the measurement of the glomerular filtration rate (GFR) as we had mentioned in our last report.³

Diabetic retinopathy is a serious sight-threatening complication of diabetes.⁴ Our teams are also formidably referring and requesting fundoscopy to screen for diabetic retinopathy. We are cognizant that although ordered, there is a challenge with getting full counter-referral response monitored and document from our service providers. This remains a challenge that NHI must solve. We look forward to reporting that this issue is solved soon.

"Must Do" Process Criteria	DAN	Hopkins	Pomona
At least annually the feet have been assessed	2.0	2.0	2.0
At least annually the patient's urine has been checked for albumin to detect early evidence of nephropathy.	2.0	2.0	2.0
At least annually the fundi have been examined for retinopathy through either examination with direct fundoscopy with dilated pupil, fundi photo or screened by ophthalmologist.	2.0	2.0	2.0

"Must Do" Process Criteria	IND	Bella Vista	Placencia	San Juan
At least annually the feet have been assessed	2.0	2.0	2.0	2.0
At least annually the patient's urine has been checked for albumin to detect early evidence of nephropathy.	2.0	2.0	2.0	2.0
At least annually the fundi have been examined for retinopathy through either examination with direct fundoscopy with dilated pupil, fundi photo or screened by ophthalmologist.	2.0	2.0	2.0	2.0

"Must Do" Process Criteria	PG	Big Falls	Santa Ana
At least annually the feet have been assessed	2.0	2.0	2.0
At least annually the patient's urine has been checked for albumin to detect early evidence of nephropathy.	2.0	2.0	2.0
At least annually the fundi have been examined for retinopathy through either examination with direct fundoscopy with dilated pupil, fundi photo or screened by ophthalmologist.	2.0	2.0	2.0

"Must Do" Process Criteria	SA	Pueblo Viejo	Santa Teresa	SPColumbia
At least annually the feet have been assessed	2.0	2.0	2.0	2.0
At least annually the patient's urine has been checked for albumin to detect early evidence of nephropathy.	2.0	2.0	2.0	2.0
At least annually the fundi have been examined for retinopathy through either examination with direct fundoscopy with dilated pupil, fundi photo or screened by ophthalmologist.	2.0	2.0	2.0	2.0

As per the tables below:

We reiterate that smoking is a known cause of type 2 diabetes. Smokers are 30–40% more likely to develop type 2 diabetes than non-smokers. And people with diabetes who smoke are more likely than non-smokers to have trouble with insulin dosing and with controlling their disease. Smokers with diabetes have higher risks for serious complications.⁵ Our clinics continue to screen our diabetic patients on the hazards of smoking and the benefits of ceasing. NHI is planning to introduce the mental health Gap Action Plan (mhGAP) developed by PAHO in 2019 to the PCPs. This tool has a stimulant dependence section from which teams may fortify their knowledge in their approach to this stimulant (nicotine) addiction.⁶⁷ This is a start.

The coexistence of hypertension in diabetes is particularly pernicious because of the strong linkage of the two conditions with all CVD, stroke, progression of renal disease and diabetic retinopathy.⁸ With this in mind our clinics are all measuring our diabetic patient's blood pressure consistently to prevent CVD events and retinopathy. All clinics scored very well here.

Our clinics are doing very well in ordering the respective lipid screens in our diabetic patients. Diabetes mellitus is considered a high- or very high-risk condition for atherosclerotic cardiovascular disease (ASCVD).⁹ Heart disease mortality is two to four times higher in patients with diabetes compared with those without diabetes. Managing the high risk for cardiovascular morbidity and mortality in diabetic patients is a challenge for practicing clinicians. Reducing the burden of cardiovascular disease in diabetes begins with assessment and treatment of elevated LDL cholesterol.¹⁰ At our clinics lipids are also being checked according to the rationalization tool for medical services.¹¹ A value added use of the lipid scores lies in its use in the atherosclerotic cardiovascular disease risk estimator plus (ASCVD risk estimator plus.) This was introduced and accepted by our clinical teams. It utilizes lipid scores, blood pressure, and medical history to "estimate our patient's 10-year ASCVD risk at an initial visit to establish a reference point."¹² Our clinics acceptance and initial use of this estimator was excellent. We continue to use tools that will promote education and advocacy for lifestyle modification on our patient's behalf.

Our clinics are doing very well identifying target organ damage and modifying treatments to these conditions. It is essential to investigate target organ damage in the diabetic. Up to 75% of adults with diabetes also have hypertension. Thus, hypertension and diabetes are common, intertwined conditions that share a significant overlap in complications. Complications include microvascular and macrovascular disorders. Macrovascular complications, which are well recognized in patients with longstanding diabetes include coronary artery disease, myocardial infarction, stroke, congestive heart failure, and peripheral vascular disease. Although microvascular complications (retinopathy, nephropathy, and neuropathy) are conventionally linked to hyperglycemia, studies have shown that hypertension constitutes an important risk factor, especially for nephropathy.¹³ Thus our viewing our patients holistically has aided in identification of these damages to target organ damage.

"Must Do" Process Criteria	DAN	Hopkins	Pomona
At least annually there has been assessment of the smoking habit	2.0	2.0	2.0
The blood pressure has been checked at every diabetes visit	2.0	2.0	2.0
At least annually the blood lipid has been checked	2.0	2.0	2.0
If any abnormal finding related to potential target organ damage was noted, the appropriate action was taken.	2.0	2.0	2.0

"Must Do" Process Criteria	IND	Bella Vista	Placencia	San Juan
At least annually there has been assessment of the smoking habit	2.0	2.0	2.0	2.0
The blood pressure has been checked at every diabetes visit	2.0	2.0	2.0	2.0
At least annually the blood lipid has been checked	2.0	2.0	2.0	2.0
If any abnormal finding related to potential target organ damage was noted, the appropriate action was taken.	2.0	2.0	2.0	2.0

"Must Do" Process Criteria	PG	Big Falls	Santa Ana
At least annually there has been assessment of the smoking habit	2.0	2.0	2.0
The blood pressure has been checked at every diabetes visit	2.0	2.0	2.0
At least annually the blood lipid has been checked	2.0	2.0	2.0
If any abnormal finding related to potential target organ damage was noted, the appropriate action was taken.	2.0	2.0	2.0

"Must Do" Process Criteria	SA	Pueblo Viejo	Santa Teresa	SPColumbia
At least annually there has been assessment of the smoking habit	2.0	2.0	2.0	2.0
The blood pressure has been checked at every diabetes visit	2.0	2.0	2.0	2.0
At least annually the blood lipid has been checked	2.0	2.0	2.0	2.0
If any abnormal finding related to potential target organ damage was noted, the appropriate action was taken.	2.0	2.0	2.0	2.0

"Should Do" Process Criteria:

These place a high emphasis on the proper follow up, education and adherence counselling offered to patients. These interventions if applied, further prevent potential target organ damage and assesses the overall outcome of the current management of the patients.¹

As per the tables below:

Our clinics are giving continues education to our patients about the importance of managing their diabetes. Although we may not have many newly diagnosed patients. The practice of reminding patients of the lifestyle recommendations is a part of the culture of treatment at our clinics.

Diabetes mellitus is a group of metabolic diseases characterized by chronic hyperglycemia.¹⁴ Caloric and national intake is a key component of diabetes management and a nutritionist can be a key stakeholder in our patients dietary education. Our referral to the nutritionist, when available for consultation at the clinics is also doing well. There has been an increase in the referral to this service. In areas, where the there is not formal nutritionis the physician reviews the DASH diet (Dietary Approaches to Stop Hypertension) guidelines with our patients and reinforces good nutritional habits. This education is applicable to diabetics and hypertensives.

A high BMI can be an indicator of high body fatness.¹⁵ Obesity and its concomitant complications (diabetes, hypertension, cardiovascular disease, and cancer, etc.) are a key component of increased risk of diabetes.¹⁶ Our NHI health teams are measuring BMI's. All the clinics are measuring BMI's more than once yearly and tracking this health indicator.

"Should Do" Process Criteria	DAN	Hopkins	Pomona
Each newly diagnosed patient has received education about diabetes management	2.0	2.0	na
At least annually the patient' s diet has been reviewed by a nutritionist with emphasis on glyceimic index	2.0	2.0	2.0
At least annually assessment of Body Mass Index has been checked	2.0	2.0	2.0

"Should Do" Process Criteria	IND	Bella Vista	Placencia	San Juan
Each newly diagnosed patient has received education about diabetes management	2.0	2.0	2.0	2.0
At least annually the patient's diet has been reviewed by a nutritionist with emphasis on glyceimic index	2.0	2.0	2.0	2.0
At least annually assessment of Body Mass Index has been checked	2.0	2.0	2.0	2.0

"Should Do" Process Criteria	PG	Big Falls	Santa Ana
Each newly diagnosed patient has received education about diabetes management	2.0	2.0	2.0
At least annually the patient's diet has been reviewed by a nutritionist with emphasis on glyceimic index	2.0	2.0	2.0
At least annually assessment of Body Mass Index has been checked	2.0	2.0	2.0

"Should Do" Process Criteria	SA	Pueblo Viejo	Santa Teresa	SPColumbia
Each newly diagnosed patient has received education about diabetes management	2.0	2.0	2.0	2.0
At least annually the patient's diet has been reviewed by a nutritionist with emphasis on glyceimic index	2.0	2.0	2.0	2.0
At least annually assessment of Body Mass Index has been checked	2.0	2.0	2.0	2.0

As per the tables below:

Creatinine is produced in muscle metabolism and excreted principally by way of the kidneys, predominantly by glomerular filtration. Blood assays for creatinine constitute the most commonly used measure of the presence and progression of CKD. Thus, the level of the serum creatinine in a subject is a general reflection of the level of kidney function. With kidney disease and loss of nephrons, the level of serum creatinine would therefore show an upward trend.¹⁷ Our clinical teams are requesting renal function (creatinine, blood urea nitrogen) regularly and homogenously to our diabetic patients. These are important in the calculation of the estimated glomerular filtration rate for the early detection of chronic kidney disease.

Adherence to medical treatment and to an improved lifestyle for the control and management of diabetes is being done homogenously and regularly at all our clinics. This function has become part of the culture of management of our diabetic population.

The annual summary form review is also being done with excellence by the clinical teams. This form assesses the general health condition of diabetics and serve a summary of the basic health indicators diabetics and their physician should have in mind in overall monitoring and treatment.¹ Our clinics have improved and are maintaining a high level on constancy in the filling out of this form.

"Should Do" Process Criteria	DAN	Hopkins	Pomona
At least annually renal function has been assessed (confirmed by BUN and CREAT)	2.0	2.0	2.0
At least annually patient has been educated on the status and importance of adherence	2.0	2.0	2.0
An annual summary review done to assess Hypertension management. Summary sheet should indicate suggested modifications proposed in the action plan, discuss lifestyle practices, symptoms and current medication regimens/and or adjustments. (review of annual summary and annual form)	2.0	2.0	2.0

"Should Do" Process Criteria	IND	Bella Vista	Placencia	San Juan
At least annually renal function has been assessed (confirmed by BUN and CREAT)	2.0	2.0	2.0	2.0
At least annually patient has been educated on the status and importance of adherence	2.0	2.0	2.0	2.0
An annual summary review done to assess Hypertension management. Summary sheet should indicate suggested modifications proposed in the action plan, discuss lifestyle practices, symptoms and current medication regimens/and or adjustments. (review of annual summary and annual form)	2.0	2.0	2.0	2.0

"Should Do" Process Criteria	PG	Big Falls	Santa Ana
At least annually renal function has been assessed (confirmed by BUN and CREAT)	2.0	2.0	2.0
At least annually patient has been educated on the status and importance of adherence	2.0	2.0	2.0
An annual summary review done to assess Hypertension management. Summary sheet should indicate suggested modifications proposed in the action plan, discuss lifestyle practices, symptoms and current medication regimens/and or adjustments. (review of annual summary and annual form)	2.0	2.0	2.0

"Should Do" Process Criteria	SA	Pueblo Viejo	Santa Teresa	SPColumbia
At least annually renal function has been assessed (confirmed by BUN and CREAT)	2.0	2.0	2.0	2.0
At least annually patient has been educated on the status and importance of adherence	2.0	2.0	2.0	2.0
An annual summary review done to assess Hypertension management. Summary sheet should indicate suggested modifications proposed in the action plan, discuss lifestyle practices, symptoms and current medication regimens/and or adjustments. (review of annual summary and annual form)	2.0	2.0	2.0	2.0

Outcome Criteria:

Since the objective of performance measures is to improve the care and health of specified populations, it makes sense to examine outcomes. Treatment of chronic conditions has shifted over the past decade to an approach that is more patient centred and that places the person with diabetes in a joint decision-making role along with the health care professionals. Patient-centered care is respectful of and responsive to individual patient preferences, needs, and values and ensures that patient values guide all clinical decisions. With this in mind, using evidence-based guidance we have set an initial set of outcome indicators. These aimed and measuring the impact of the health interventions in mitigating the negative impact of diabetes.¹

As per the tables below:

HbA1c is challenging to control in the general diabetic patient. The auditing team considered this great difficulty. Our consultant internist agreed that in objective fairness to our clinical teams that we should look to see if an HbA1c baseline was sought to be established and if so, a 2-point score was given. Also, if in the follow-up HbA1c levels found, the 7% is not attainable at the time of assessments, due to challenges not currently within the control of the PCP, that a full score would be given. We note that all our PCPs are following the medical treatment and follow-up guidelines that are nationally and internationally recognized in terms of HbA1c.

Our audit and management team are now planning to look at these outcome measure definitions to redefine their expected goals and the way we audit them. This is to aid in continued objectivity in the audit while being fair to the PCPs in their engaging patients to lower and maintain control of their HbA1c.

Outcome Criteria	DAN	Hopkins	Pomona
Patient's Hb A1c target established is appropriate if the 7% is not attainable at the time of assessment	2.0	2.0	2.0
Patient's Hb A1c target achieved	2.0	2.0	2

Outcome Criteria	IND	Bella Vista	Placencia	San Juan
Patient's Hb A1c target established is appropriate if the 7% is not attainable at the time of assessment	2.0	2.0	2.0	2.0
Patient's Hb A1c target achieved	2.0	2.0	2.0	2.0

Outcome Criteria	PG	Big Falls	Santa Ana
Patient's Hb A1c target established is appropriate if the 7% is not attainable at the time of assessment	2.0	2.0	2.0
Patient's Hb A1c target achieved	2.0	2.0	2.0

Outcome Criteria	SA	Pueblo Viejo	Santa Teresa	SPColumbia
Patient's Hb A1c target established is appropriate if the 7% is not attainable at the time of assessment	2.0	2.0	2.0	2.0
Patient's Hb A1c target achieved	2.0	2.0	2.0	2.0

Blood Pressure Monitoring:

Age >= 60 with BP <150/90 * PCP Crosstabulation

			PCP				Total
			DAN	IND	PG	SA	
>60 150/90	Partially met	Count	1	0	0	0	1
		% within PCP	3.4%	.0%	.0%	.0%	.8%
	Fully met	Count	6	24	18	28	76
		% within PCP	20.7%	70.6%	62.1%	82.4%	60.3%
	NA	Count	22	10	11	6	49
		% within PCP	75.9%	29.4%	37.9%	17.6%	38.9%
	Total	Count	29	34	29	34	126
		% within PCP	100.0%	100.0%	100.0%	100.0%	100.0%

Our teams are consistently following guidelines measuring our diabetic patient's blood pressure in our >60 years patients. This ensures that any patient that is hypertensive will be monitored and get concomitant treatment and guidance in this regard. The partially met criteria are for patients who had spikes of >150 / > 90 of blood pressure but maintained the suggested scores more constantly. Most patients who are > 60 years have fully met the criteria by having a blood pressure <150/<90.

Ages >18 to 59 yrs, BP <140/<90 * PCP Crosstabulation

			PCP				Total
			DAN	IND	PG	SA	
>18 140/90	Fully met	Count	20	10	11	6	47
		% within PCP	69.0%	29.4%	37.9%	17.6%	37.3%
	NA	Count	9	24	18	28	79
		% within PCP	31.0%	70.6%	62.1%	82.4%	62.7%
Total		Count	29	34	29	34	126
		% within PCP	100.0%	100.0%	100.0%	100.0%	100.0%

Our teams are consistently following guidelines measuring our diabetic patient's blood pressure in our 18-59 years patients. This ensures that any patient that is hypertensive will be monitored and get concomitant treatment and guidance in this regard. The partially met criteria are for patients who had spikes of >140 / > 90 of blood pressure but maintained the suggested scores more constantly. Most patients who are 18-59 years have fully met the criteria by having a blood pressure <140/<90.

As per the tables below:

Low-density lipoprotein (LDL) is being measured at all our clinics in a consistent manner. This action has enabled for the screening of atherosclerotic cardiovascular heart disease risk estimation to be adopted, since we need the LDL and other lipid scores for risk estimation.

Outcome Criteria	DAN	Hopkins	Pomona
Patient's LDL cholesterol level is less than 135 mg/dl	2.0	2.0	2

Outcome Criteria	IND	Bella Vista	Placencia	San Juan
Patient's LDL cholesterol level is less than 135 mg/dl	2.0	2.0	2.0	2.0

Outcome Criteria	PG	Big Falls	Santa Ana
Patient's LDL cholesterol level is less than 135 mg/dl	2.0	2.0	2.0

Hypertension Audit Results 2018-2019:

2018 – 2019 Overall Scores	DAN	IND	PG	SA
Scores (target 90%)	100%	100%	100%	100%

All PCP clinics scored over their target of 90%. This signifies that have established a very good baseline standard of care. This is homogenized at all the PCPs. Dangriga (2017-2018: 90%), Independence (2017-2018: 91%), Punta Gorda (2017-2018: 90%) and San Antonio (2017-2018: 88%) have all had improvement in respect of last year's clinical audit. Our clinics are to all be commended for these results. The records that the clinical teams are keeping are being done very well.

	DAN	Hopkins	Pomona	Score
# of records assessed	19	5	5	29
# of records that met the target (90%)	19	5	5	29
Total score	100%	100%	100%	100%

	IND	Bella Vista	Placencia	San Juan	Score
# of records assessed	19	5	5	5	34
# of records that met the target (90%)	19	5	5	5	34
Total score	100%	100%	100%	100%	100%

	PG	Big Falls	Santa Ana	Score
# of records assessed	19	5	5	29
# of records that met the target (90%)	19	5	5	29
Total score	100%	100%	100%	100%

	SA	Pueblo Viejo	Santa Teresa	SPColumbia	Score
# of records assessed	19	5	5	5	34
# of records that met the target (90%)	19	5	5	5	34
Total score	100%	100%	100%	100%	100%

Demographics	DAN	IND	PG	SA
Gender				
Male	11	9	3	6
	38%	27%	10%	18%
Female	18	25	26	28
	62%	74%	90%	82%

As is the usual case, the majority of people seeking medical attention are women. Overall 77% of the audited files were female, versus 23% for males. Punta Gorda had 90% of their patients being women. Primary care would need to develop an improved value proposition for our male patients, to improve their visitation for their healthcare needs.

Demographics	DAN	IND	PG	SA
Age Range				
31-45	0	7	8	3
	0%	21%	28%	9%
46-55	9	10	2	7
	31%	29%	7%	21%
>55	20	16	19	24
	79%	50%	41%	68%

The majority of patients that are currently being treated for hypertension are >55 years-old for all the clinics. On averages 60% of our patients are >55 years/old; this is the same for the 2017-2018 clinical audit. It is concerning that 36% of our overall patients are between 31-55 years-old. Dangriga has 31%, Independence 50%, Punta Gorda 35% and San Antonio 30% of their patient cohort is in this age range.

Demographics	DAN	IND	PG	SA
Ethnicity				
Creole	3	11	3	1
	10%	32%	10%	3%
Mestizo	8	16	8	1
	28%	47%	28%	3%
Garifuna	18	3	2	0
	62%	9%	7%	0%
Maya	0	0	7	26
	0%	0%	24%	77%

The predominant ethnic groups in the southern clinics are Mestizo (overall average of 26% of the population) and Maya (25%). Garinagu accounted for 19% of the overall population, with Dangriga having the highest overall percentage of this patients being Garinagu (62%). Independence highest demographic is Mestizo (47% of the patient charts seen). The highest prevalence of Maya was at San Antonio with 77%.

Demographics	DAN	IND	PG	SA
Case				
Newly Diagnosed	0	0	0	0
	0%	0%	0%	0%
Established Cases	27	34	28	34
	93%	100%	97%	100%

The relationship between blood pressure and risk of CVD events is continuous, consistent, and independent of other risk factors. The higher the BP, the greater the chance of heart attack, HF, stroke, and kidney diseases⁸. Hypertension requires a multifaceted approach in control. A subset of patients requires intense multidisciplinary interventions that work in conjunction with the medical therapy. Overall all cases that were being treated were already established cases. This was similar to last year's averages.

Demographics	DAN	IND	PG	SA
Classification				
Hypertension Stage 1	18	24	24	26
	62%	71%	83%	77%
Hypertension Stage 2	10	10	5	8
	35%	29%	17%	24%
Hypertension Controlled				
Yes	22	25	25	32
	76%	74%	86%	94%
No	6	9	4	2
	21%	27%	14%	6%

Hypertension targets are set by the Seventh Report of the Joint National Committee (JNC7) on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure definitions for hypertension⁸. JNC 7 suggests that all people with hypertension (stages 1 (Systolic 140-159mmHg/Diastolic 90-99mmHg) and 2 (Systolic equal or greater than 160mmHg/ Diastolic equal or greater than 100 mmHg) be treated. The treatment goal for individuals with hypertension and no other compelling conditions is <140/90 mmHg.

73% of the overall population of hypertensives are stage 1 (compared to 2017-2018 audit of: 74%) and 26% of the population are in stage 2 hypertension (2017: 16%). There is a noted increase in the percentage of stage 2 hypertensives with Dangriga and Independence having the higher number of hypertensives in their populations. PG and SA had the highest number of stage 1 hypertensives.

There was on overall increase in the number of controlled cases: 2017-2018 had 77%, and the 2018-2019 audit shows 82%. San Antonio and Punta Gorda had the highest number of controlled hypertension cases, and the highest increases since last year.

“Must Do” Criteria:

As per the tables below:

In all clinics blood pressure readings are being corroborated at second and even third instances when there are high measurements.

Hypertensive emergencies are characterized by severe elevations in blood pressure (BP >180/120 mmHg) complicated by evidence of impending or progressive target organ dysfunction. They require immediate BP reduction (not necessarily to normal) to prevent or limit target organ damage. Examples include hypertensive encephalopathy, intracerebral haemorrhage, acute myocardial infarction (MI), acute left ventricular failure with pulmonary edema, unstable angina pectoris, dissecting aortic aneurysm, or eclampsia.⁸ *Hypertensive urgencies* are those situations associated with severe elevations in BP without progressive target organ dysfunction. Examples include upper levels of stage II hypertension associated with severe headache, shortness of breath, epistaxis, or severe anxiety. The majority of these patients present as noncompliant or inadequately treated hypertensive individuals, often with little or no evidence of target organ damage.⁸ Our clinics are monitoring for hypertensive emergencies and urgencies that would require acute medical treatment. The auditors noted no patient files with hypertensive emergencies. Noted were a few with hypertensive urgencies, that were treated.

As we have highlighted previously we can note that hypertension classification based on JNC 7 is being done consistently well throughout our program. JNC 7 suggests that all people with hypertension (stages 1 (Systolic 140-159mmHg/Diastolic 90-99mmHg) and 2 (Systolic equal or greater than 160mmHg/ Diastolic equal or greater than 100 mmHg) be treated. The treatment goal for individuals with hypertension and no other compelling conditions is <140/90 mmHg.⁸ All our patients were well classified according to JNC 7 and were in their appropriate classification.

Must Do Criteria			
Diagnosing hypertension/Follow up	DAN	Hopkins	Pomona
If blood pressure measured in the clinic is 140/90 mmHg or higher: <ul style="list-style-type: none">• A second measurement during the consultation was taken.• If the second measurement was substantially different from the first, a third measurement was taken.	2.0	2.0	2.0
If the person presented with severe hypertension, anti-hypertensive drug treatment therapy was administered immediately, without waiting for the results of ABPM or HBPM.	2.0	1.9	NA
Hypertension Classification was appropriate based on subsequent BPM readings. (at least three consecutive readings on different consultation days)	2.0	2.0	2.0

Must Do Criteria				
Diagnosing hypertension/Follow up	IND	Bella Vista	Placencia	San Juan
<p>If blood pressure measured in the clinic is 140/90 mmHg or higher:</p> <ul style="list-style-type: none"> • A second measurement during the consultation was taken. • If the second measurement was substantially different from the first, a third measurement was taken. 	2.0	2.0	2.0	2.0
If the person presented with severe hypertension, anti-hypertensive drug treatment therapy was administered immediately, without waiting for the results of ABPM or HBPM.	2.0	2.0	2.0	2.0
Hypertension Classification was appropriate based on subsequent BPM readings. (at least three consecutive readings on different consultation days)	2.0	2.0	2.0	2.0

Must Do Criteria			
Diagnosing hypertension/Follow up	PG	Big Falls	Santa Ana
<p>If blood pressure measured in the clinic is 140/90 mmHg or higher:</p> <ul style="list-style-type: none"> • A second measurement during the consultation was taken. • If the second measurement was substantially different from the first, a third measurement was taken. 	2.0	2.0	2.0
If the person presented with severe hypertension, anti-hypertensive drug treatment therapy was administered immediately, without waiting for the results of ABPM or HBPM.	NA	2.0	2.0
Hypertension Classification was appropriate based on subsequent BPM readings. (at least three consecutive readings on different consultation days)	2.0	2.0	2.0

Must Do Criteria				
Diagnosing hypertension/Follow up	SA	Pueblo Viejo	Santa Teresa	SPColumbia
<p>If blood pressure measured in the clinic is 140/90 mmHg or higher:</p> <ul style="list-style-type: none"> • A second measurement during the consultation was taken. • If the second measurement was substantially different from the first, a third measurement was taken. 	2.0	2.0	2.0	2.0
If the person presented with severe hypertension, anti-hypertensive drug treatment therapy was administered immediately, without waiting for the results of ABPM or HBPM.	2.0	2.0	2.0	NA
Hypertension Classification was appropriate based on subsequent BPM readings. (at least three consecutive readings on different consultation days)	2.0	2.0	2.0	2.0

As per the tables below:

In the assessment of cardiovascular damage and target organ damage our clinics are monitoring the target organs. This is crucial for patient management. As mentioned before, the relationship between blood pressure and risk of cardiovascular disease (CVD) events is continuous, consistent, and independent of other risk factors. The higher the BP, the greater the chance of heart attack, HF, stroke, and kidney diseases.⁸ Hypertensive patients are at increased risk for myocardial infarction (MI) or other major coronary events and may be at higher risk of death following an acute MI. Cardiomegaly in these hypertensive cases is usually due to left ventricular hypertrophy (LVH), which is an increase in left ventricular mass. In untreated or poorly treated individuals, LVH becomes a major risk factor for dilated cardiomyopathy and heart failure (HF). The coexistence of hypertension in diabetes is particularly pernicious because of the strong linkage of the two conditions with all CVD, stroke, progression of renal disease and diabetic retinopathy.⁸

Our clinics are screening patients for other target organ damage: chronic heart failure, angina, stroke history, transient ischemic attack, hypertensive kidney disease and peripheral artery disease. Electrocardiograms are being done and the internist noted that the ones reviewed are technically correct. There is sporadic discrepancy in its interpretation, but this is controlled, particularly with internal medicine referrals.

Of note dementia and cognitive impairment occur more commonly in people with hypertension. Reduced progression of cognitive impairment may occur with effective antihypertensive therapy.⁸ As our population ages and lives longer we will need to improve our monitoring of the effects of hypertension on brain function. Our clinics are screening patients very well for the stroke and transient ischemic attack history.

Cardiovascular disease (CVD) is the most common cause of death in individuals with CKD, and CKD is an independent risk factor for CVD. Microalbuminuria (30–300 mg/day) is associated with increased CVD risk in diabetics and other high-risk patients. Urinary albumin excretion has diagnostic and prognostic value equivalent to reduced estimated glomerular filtration rates (eGFRs).⁸ Creatinine and blood urea nitrogen are being monitored in patients consistently, and this is important in order to calculate the GFR. Our internal medicine auditors congratulate NHI and their teams for monitoring of the glomerular filtration rate (GFR).³ This would be another layer in outcome measurement that would add more detail to the renal function of patients. This was a recommendation from last years clinical audit that has come to fruition.

Peripheral artery disease (PAD) is a narrowing of the peripheral arteries to the legs, stomach, arms, and head — most commonly in the arteries of the legs. PAD is similar to coronary artery disease (CAD). Both PAD and CAD are caused by atherosclerosis that narrows and blocks arteries in various critical regions of the body.¹⁸ Major risk factors for peripheral arterial disease (PAD) are hypertension, diabetes, and smoking. Symptomatic PAD is associated with a greatly increased risk of death from CVD, in part because diffuse atherosclerosis, CAD, and renovascular disease frequently coexist in these patients. Therefore, more intensive screening for these related cardiovascular disorders is appropriate in persons with PAD.⁸ Our clinics are documenting the search for PAD.

Differences in systolic blood pressure (SBP) of 10 mm Hg or more or 15 mm Hg or more between arms have been associated with peripheral vascular disease and attributed to subclavian stenosis. It is a useful indicator of risk of vascular disease and death.¹⁹ Our clinics are screening well for vascular disease compromise, as they all scored 2 in this area.

Hypertension can affect the retina, choroid, and optic nerve of the eye, particularly with stage 2 hypertension. These changes can be appreciated with inspection of the retinal vessels by direct ophthalmoscopy.⁸ Our clinics are complying with these screening guidelines and ordering these for our hypertensive patients consistently.

The auditors noted that there is a still a gap in ordering ophthalmology screens and receiving a result report at this clinic for physician reference. This a process that NHI shall endeavour to fix.

Target organ damage is being sought out for diagnosis and prevention in our patients. The auditing team noted that our PCPs and satellites in the SR performed appropriately in target organ damage and took appropriate action in these cases.

Must Do Criteria			
Assessing cardiovascular risk and target organ damage	DAN	Hopkins	Pomona
The records show that at diagnosis the following symptoms and signs of target organ damage have been sought:			
HEART:	1.9	1.6	2.0
LVH/Myocardial Infarction: EKG			
Chronic Heart Failure: EKG; Chest Xray			
Angina: Reported history			
BRAIN:	2.0	2.0	2.0
Strokes: Reported history			
Transient Ischemic Attack: Reported History			
Kidney:	2.0	1.92	2.0
Proteinuria/Microalbuminuria: Lab test results			
BUN/Creatinine			
Vascular Disease:	2.0	2.0	2.0
Peripheral Arterial disease: Patient History			
Assymetrical Pulses: BPM in both arms and EKG			
Eyes:	2.0	2.0	2.0
Retinopathy: eye fundi exam			
If any abnormal finding related to potential target organ damage was noted, the appropriate action was taken.	2.0	2.0	2.0

Must Do Criteria				
Assessing cardiovascular risk and target organ damage	IND	Bella Vista	Placencia	San Juan
The records show that at diagnosis the following symptoms and signs of target organ damage have been sought:				
HEART:	2.0	2.0	2.0	2.0
LVH/Myocardial Infarction: EKG				
Chronic Heart Failure: EKG; Chest Xray				
Angina: Reported history				
BRAIN:	2.0	2.0	2.0	2.0
Strokes: Reported history				
Transient Ischemic Attack: Reported History				
Kidney:	2.0	2.0	2.0	2.0
Proteinuria/Microalbuminuria: Lab test results				
BUN/Creatinine				
Vascular Disease:	2.0	2.0	2.0	2.0
Peripheral Arterial disease: Patient History				
Assymetrical Pulses: BPM in both arms and EKG				
Eyes:	2.0	2.0	2.0	2.0
Retinopathy: eye fundi exam				
If any abnormal finding related to potential target organ damage was noted, the appropriate action was taken.	2.0	2.0	2.0	2.0

Must Do Criteria			
Assessing cardiovascular risk and target organ damage	PG	Big Falls	Santa Ana
The records show that at diagnosis the following symptoms and signs of target organ damage have been sought:			
HEART:	2.0	2.0	2.0
LVH/Myocardial Infarction: EKG			
Chronic Heart Failure: EKG; Chest Xray			
Angina: Reported history			
BRAIN:	2.0	2.0	2.0
Strokes: Reported history			
Transient Ischemic Attack: Reported History			
Kidney:	2.0	2.0	2.0
Proteinuria/Microalbuminuria: Lab test results			
BUN/Creatinine			
Vascular Disease:	2.0	2.0	2.0
Peripheral Arterial disease: Patient History			
Assymetrical Pulses: BPM in both arms and EKG			
Eyes:	1.9	2.0	2.0

Retinopathy: eye fundi exam			
If any abnormal finding related to potential target organ damage was noted, the appropriate action was taken.	2.0	2.0	2.0

Must Do Criteria				
Assessing cardiovascular risk and target organ damage	SA	Pueblo Viejo	Santa Teresa	SPColumbia
The records show that at diagnosis the following symptoms and signs of target organ damage have been sought:				
HEART:	2.0	2.0	2.0	2.0
LVH/Myocardial Infarction: EKG				
Chronic Heart Failure: EKG; Chest Xray				
Angina: Reported history				
BRAIN:	2.0	2.0	2.0	2.0
Strokes: Reported history				
Transient Ischemic Attack: Reported History				
Kidney:	2.0	2.0	2.0	2.0
Proteinuria/Microalbuminuria: Lab test results				
BUN/Creatinine				
Vascular Disease:	2.0	2.0	2.0	2.0
Peripheral Arterial disease: Patient History				
Assymetrical Pulses: BPM in both arms and EKG				
Eyes:	2.0	2.0	2.0	2.0
Retinopathy: eye fundi exam				
If any abnormal finding related to potential target organ damage was noted, the appropriate action was taken.	2.0	2.0	2.0	2.0

“Should do” Criteria:

These place a high emphasis on the proper follow up, education and adherence counselling offered to patients. These interventions if applied, further prevent potential target organ damage and assesses the overall outcome of the current management of the patients.¹

As per the tables below:

The benefits of modest **lifestyle** changes on cardiovascular risk factors are well documented. Behavioural models suggest that the most effective therapy prescribed by the most careful clinician will control hypertension only if the patient is motivated to take the medication as directed and to establish and maintain a health-promoting lifestyle. Motivation improves when patients have positive experiences with, and trust in, their clinicians. Better communication improves outcomes; empathy builds trust and is a potent motivator.⁸

Comparing these results to our last clinics audit we note continued good practices with all our clinics scoring high here. Overall our patients are being spoken to and guided by our clinical teams in lifestyle modifications for the improvement of their hypertension. Our clinics are documenting their strategies in lifestyle modification and have scored well in this regard. This continues to be the bedrock of our public health treatment strategy.

The emphasis for **weight management** should be on avoidance of excess total energy intake and a regular pattern of physical activity. Reducing food portion sizes and limiting fat intake can assist in reducing overall calorie intake. For example, adoption of the well-studied low sodium DASH eating plan provides heart healthy foods that can be used to promote weight loss, reduce blood pressure in both hypertensive and prehypertensive individuals, and reduce low density lipoproteins (LDL).⁸

Adoption of the well-studied **low sodium** DASH (Dietary Approaches to Stop Hypertension) eating plan provides heart healthy foods that can be used to promote weight loss, reduce blood pressure in both hypertensive and prehypertensive individuals, and reduce low density lipoproteins (LDL).⁸ Our clinics continue to promote this diet and speaking to patients about the realities of their dietary needs, tailoring the diets to the needs of patients. It is important for a nutritionist to be available to continue to lay good dietary foundations for our hypertensive population.

Increased **physical activity**, when combined with a reduction in calories, is essential to weight loss success. Based on the available evidence, the recommendation is to engage in regular physical activity at least 30 minutes per day, most days of the week (see table 9). In addition, physical activity is critical to the maintenance of weight loss and is important for overall reduction in cardiovascular risk; 60–90 minutes per week of walking can reduce CHD mortality by about 50 percent.⁸

NHI does recognize that we need to embark on innovative strategies and revisit past ones to improve patient buy-in for lifestyle interventions adoption.

Consumption of **alcohol** (e.g., >30 grams of ethanol a day or approximately two “drinks” daily) have a dose-related effect on BP, both in hypertensive and normotensive subjects.⁸ Our at-risk patients received information to prevent the excessive use of alcohol intake in all our clinics.

Physicians should be aware of the possible indications and side effects related to the use of **herbal medications**.

Physicians should improve their knowledge of herbal medications to adequately weigh the clinical implications related to their use, and be able to discuss with patients their possible benefits and side effects, and explain that natural does not always mean safe.²⁰

Use of herbal medications for the treatment of cardiovascular diseases is not supported by scientific evidence. Although most of the herbs demonstrate an effect on biological mechanisms associated with cardiovascular disease, available clinical studies are limited in sample size and appropriate controls. Physicians should be certain patients inform them regarding all supplements, particularly herbal, which can result in serious bleeding and drug interactions. And physicians should try to assess whether patients are not taking their medication as prescribed because of the preference for herbal supplements, which they perceive as safe and not “drugs.”²¹

However, correctly identifying new signs or symptoms of herbal medications is challenging for physicians, due to a lack of knowledge on their effects and common under-reporting from patients. Indeed, about 70% of patients do not notify their physicians about their use of supplements and, at the same time, physicians are unlikely to regularly gather correct information on their use. In fact, on the one hand patients believe that it is pointless to report the use of herbal medications, because they are reputed to be safe and are not perceived as drugs, whereas on the other hand, physicians do not give adequate weight to the clinical implications of such preparations. Moreover, use of herbal medications has been associated with poor adherence to conventional medications, raising significant concerns in terms of safety and effectiveness of concurrent treatments.²⁰

To deal with this issue, physicians should carry out a structured interview. Information on use of alternative preparations, herbal medications, and actual drugs should be properly recorded, and signs and symptoms must be evaluated in accordance with the clinical story. Communicating with the patient is a crucial component of the process, where the pros and cons of specific herbal medications should be explained and their risk–benefit profile properly discussed. Eventually, the decision to withdraw or withhold a new herbal medication should be shared with the patient.²⁰

Our clinics are discussing the use of alternative medications. This question is of continued relevance with the advent of commercialization and the social media bombardment of patients with multi-cure medications and natural cures. These alternative treatments pose a real barrier to medication adherence and a communication strategy for the bilateral understanding of their use and misuse needs to be developed.

Smoking raises blood pressure acutely. Smoking cessation may be the single most important factor whether peripheral artery disease (PAD) progresses. For overall cardiovascular risk reduction, patients should be strongly counselled to quit smoking.⁸ We also know smoking causes type 2 diabetes. In fact, smokers are 30–40% more likely to develop type 2 diabetes than non-smokers. And people with diabetes who smoke are more likely than non-smokers to have trouble with insulin dosing and with controlling their disease. Smokers with diabetes have higher risks for serious complications.⁵

“Effective tobacco dependence treatment should include cessation advice incorporated into primary health-care services... and access to free or low-cost cessation medicines. In Belize there is no smoking cessation support available. Some pharmacological treatment is available.”²² Our clinical teams continue to screen the population for tobacco smoking. Patients in general replied that they did not at all clinics. Our internal medicine auditor makes note that treatment need to be addressed for smokers.

Should Do Criteria			
Lifestyle interventions	DAN	Hopkins	Pomona
Lifestyle advice was offered initially and then periodically to people undergoing assessment or treatment for hypertension.	2.0	2.0	2.0
Appropriate guidance on diet and exercise as part of the action plan was offered.	2.0	2.0	2.0
At least annually, alcohol consumption was assessed and advice on reduced intake offered.	2.0	2.0	2.0
Advice on dietary sodium intake offered, (either by reducing or substituting sodium salt and educating patients on hidden sources of sodium)	2.0	2.0	2.0
Discuss alternative medicines used by patients for reducing blood pressure.	2.0	2.0	2.0
At least annually smoking practice was assessed and cessation advice offered.	2.0	2.0	2.0

Should Do Criteria				
Lifestyle interventions	IND	Bella Vista	Placencia	San Juan
Lifestyle advice was offered initially and then periodically to people undergoing assessment or treatment for hypertension.	2.0	2.0	2.0	2.0
Appropriate guidance on diet and exercise as part of the action plan was offered.	2.0	2.0	2.0	2.0
At least annually, alcohol consumption was assessed and advice on reduced intake offered.	2.0	2.0	2.0	2.0
Advice on dietary sodium intake offered, (either by reducing or substituting sodium salt and educating patients on hidden sources of sodium)	2.0	2.0	2.0	2.0
Discuss alternative medicines used by patients for reducing blood pressure.	2.0	2.0	2.0	2.0
At least annually smoking practice was assessed and cessation advice offered.	2.0	2.0	2.0	2.0

Should Do Criteria

Lifestyle interventions	PG	Big Falls	Santa Ana
Lifestyle advice was offered initially and then periodically to people undergoing assessment or treatment for hypertension.	2.0	2.0	2.0
Appropriate guidance on diet and exercise as part of the action plan was offered.	2.0	2.0	2.0
At least annually, alcohol consumption was assessed and advice on reduced intake offered.	2.0	2.0	2.0
Advice on dietary sodium intake offered, (either by reducing or substituting sodium salt and educating patients on hidden sources of sodium)	2.0	2.0	2.0
Discuss alternative medicines used by patients for reducing blood pressure.	2.0	2.0	2.0
At least annually smoking practice was assessed and cessation advice offered.	2.0	2.0	2.0

Should Do Criteria

Lifestyle interventions	SA	Pueblo Viejo	Santa Teresa	SPColumbia
Lifestyle advice was offered initially and then periodically to people undergoing assessment or treatment for hypertension.	2.0	2.0	2.0	2.0
Appropriate guidance on diet and exercise as part of the action plan was offered.	2.0	2.0	2.0	2.0
At least annually, alcohol consumption was assessed and advice on reduced intake offered.	2.0	2.0	2.0	2.0
Advice on dietary sodium intake offered, (either by reducing or substituting sodium salt and educating patients on hidden sources of sodium)	2.0	2.0	2.0	2.0
Discuss alternative medicines used by patients for reducing blood pressure.	2.0	2.0	2.0	2.0
At least annually smoking practice was assessed and cessation advice offered.	2.0	2.0	2.0	2.0

As per the tables below:

Our clinics are initiating and monitoring antihypertensive drug treatments as per international and NHI guidelines in both stages 1 and 2. They are also taking into consideration comorbidities in the treatment regimens. Our internal medicine auditor agrees that overall the treatments are appropriate.

Should Do Criteria			
Initiating and monitoring anti-hypertensive drug treatment, including blood pressure targets	DAN	Hopkins	Pomona
Initiating treatment			
Anti-hypertensive drug treatment offered to people aged under 80 years with stage 1 hypertension who have one or more of the following: <ul style="list-style-type: none"> • target organ damage • established cardiovascular disease • renal disease • diabetes 	2.0	2.0	2.0
Anti-hypertensive drug treatment to people of any age with stage 2 hypertension.	2.0	2.0	NA

Should Do Criteria				
Initiating and monitoring anti-hypertensive drug treatment, including blood pressure targets	IND	Bella Vista	Placencia	San Juan
Initiating treatment				
Anti-hypertensive drug treatment offered to people aged under 80 years with stage 1 hypertension who have one or more of the following: <ul style="list-style-type: none"> • target organ damage • established cardiovascular disease • renal disease • diabetes 	2.0	2.0	2.0	2.0
Anti-hypertensive drug treatment to people of any age with stage 2 hypertension.	2.0	2.0	2.0	2.0

Should Do Criteria

Initiating and monitoring anti-hypertensive drug treatment, including blood pressure targets	PG	Big Falls	Santa Ana
Initiating treatment			
Anti-hypertensive drug treatment offered to people aged under 80 years with stage 1 hypertension who have one or more of the following: <ul style="list-style-type: none"> • target organ damage • established cardiovascular disease • renal disease • diabetes 	2.0	2.0	2.0
Anti-hypertensive drug treatment to people of any age with stage 2 hypertension.	2.0	2.0	2.0

Should Do Criteria

Initiating and monitoring anti-hypertensive drug treatment, including blood pressure targets	SA	Pueblo Viejo	Santa Teresa	SPColumbia
Initiating treatment				
Anti-hypertensive drug treatment offered to people aged under 80 years with stage 1 hypertension who have one or more of the following: <ul style="list-style-type: none"> • target organ damage • established cardiovascular disease • renal disease • diabetes 	2.0	2.0	2.0	2.0
Anti-hypertensive drug treatment to people of any age with stage 2 hypertension.	2.0	2.0	2.0	NA

As per the tables below:

Clinical inertia is the failure to titrate or combine medications and to reinforce lifestyle modifications, despite knowing that the patient is not at goal BP. This may be due in part to clinician focus on relieving symptoms, a lack of familiarity with clinical guidelines, or dis-comfort in titrating to a goal.²³ In the majority of our patients, the treatment regimens are well met in accordance to their co-morbidities. This is also in coherence with our results of this KPI last year.

Medication side-effects are always a present risk for those on medication. Racial differences in the incidence of antihypertensive drug side effects may occur; African Americans and Asians have a three to fourfold higher risk of angioedema and have more cough attributed to ACEIs than Caucasians.²³ Our patients are being monitored for potential side-effects and we commend our clinics in continuing this target. This makes our patients safer and engenders trust from our patients that we are monitoring this. Continued communication with our patient is a corner stone of good treatment of our patients.

This summary review is important for the referencing of patient's illnesses. It enables quick reference of illness and the validation of treatment regimens in a new user of the chart. There were annual summary review forms are well filled out by our clinics as seen above. This is a crucial part of the clinical documentation for the review of treatment and the follow-up of that treatment.

Patients are being monitored for side-effects and the benefits of their treatment. Adherence is also being monitored at all clinics.

Behaviour models suggest that the most effective therapy prescribed by the most careful clinician will control hypertension only if the patient is motivated to take the medication as directed and to establish and maintain a health-promoting lifestyle. Motivation improves when patients have positive experiences with, and trust in, their clinicians. Better communication improves outcomes; empathy builds trust and is a potent motivator.⁸ It is challenging to get patient buy-in to decrease non-adherence so it's monitoring is important. All our clinics are reviewing nonadherence in high percentages with their patients. This is a crucial part of the clinical assessment and will enable clinicians to optimize treatment and get to their desired clinical outcomes.

Should Do Criteria			
Choosing antihypertensive drug treatment	DAN	Hopkins	Pomona
Treatment regimen appropriate and in accordance to reported co-morbidities	2.0	2.0	2.0
Patient education and adherence to treatment			
At least annually patient is educated on the benefits of treatment regimen and information provided on potential side effects.	2.0	2.0	2.0
Patients monitored for the presence of unwanted side effects and action taken was appropriate	1.9	2.0	2.0
An annual summary review done to assess Hypertension management. Summary sheet should indicate suggested modifications proposed in the action plan, discuss lifestyle practices, symptoms and current medication regimens/and or adjustments. (review of annual summary and annual form)	2.0	2.0	2.0
Review and document potential causes of non-adherence and respond accordingly.	2.0	2.0	2.0

Should Do Criteria				
Choosing antihypertensive drug treatment	IND	Bella Vista	Placencia	San Juan
Treatment regimen appropriate and in accordance to reported co-morbidities	2.0	2.0	2.0	2.0
Patient education and adherence to treatment				
At least annually patient is educated on the benefits of treatment regimen and information provided on potential side effects.	2.0	2.0	2.0	2.0
Patients monitored for the presence of unwanted side effects and action taken was appropriate	2.0	2.0	2.0	2.0
An annual summary review done to assess Hypertension management. Summary sheet should indicate suggested modifications proposed in the action plan, discuss lifestyle practices, symptoms and current medication regimens/and or adjustments. (review of annual summary and annual form)	2.0	2.0	2.0	2.0
Review and document potential causes of non-adherence and respond accordingly.	2.0	2.0	2.0	2.0

Should Do Criteria			
Choosing antihypertensive drug treatment	PG	Big Falls	Santa Ana
Treatment regimen appropriate and in accordance to reported co-morbidities	2.0	2.0	2.0
Patient education and adherence to treatment			
At least annually patient is educated on the benefits of treatment regimen and information provided on potential side effects.	2.0	2.0	2.0
Patients monitored for the presence of unwanted side effects and action taken was appropriate	2.0	2.0	2.0
An annual summary review done to assess Hypertension management. Summary sheet should indicate suggested modifications proposed in the action plan, discuss lifestyle practices, symptoms and current medication regimens/and or adjustments. (review of annual summary and annual form)	2.0	2.0	2.0
Review and document potential causes of non-adherence and respond accordingly.	2.0	2.0	2.0

Should Do Criteria				
Choosing antihypertensive drug treatment	SA	Pueblo Viejo	Santa Teresa	SPColumbia
Treatment regimen appropriate and in accordance to reported co-morbidities	2.0	2.0	2.0	2.0
Patient education and adherence to treatment				
At least annually patient is educated on the benefits of treatment regimen and information provided on potential side effects.	2.0	2.0	2.0	2.0
Patients monitored for the presence of unwanted side effects and action taken was appropriate	2.0	2.0	2.0	2.0
An annual summary review done to assess Hypertension management. Summary sheet should indicate suggested modifications proposed in the action plan, discuss lifestyle practices, symptoms and current medication regimens/and or adjustments. (review of annual summary and annual form)	2.0	2.0	2.0	2.0
Review and document potential causes of non-adherence and respond accordingly.	2.0	2.0	2.0	2.0

Hypertension: By age range:

Achieving blood pressure targets is challenging. Our auditors noted that in a diverse population with comorbidities it can be difficult to get to treatment goals. Partial scores were given for cases in which there was overall control of blood pressure. Our internal medicine auditor notes that referral to a specialist might be appropriate in these cases for possible scaling up of treatment.

In the patient cohort <60 yrs. old there was fully met outcome in most patients at all the clinics. There were partially met cases at IHC. Overall the management and treatment of patients <60 years old is appropriate and good. In the cases of partially met, our internal medicine auditor suggests that there be continued referral and inter-consultation with and internist to aid in getting these patient's blood pressure controlled better.

Ages >18 to 59 yrs., BP <140/<90 * PCP Crosstabulation

			Clinic				Total
			DAN	IND	PG	SA	
Score	Fully met	Count	12	22	13	15	62
		% within Clinic	100.0%	100.0%	100.0%	100.0%	100.0%
Total		Count	12	22	13	15	62
		% within Clinic	100.0%	100.0%	100.0%	100.0%	100.0%

In the patient cohort >60 yrs. old most patients met the “fully met” criteria, indicating their blood pressure was well controlled. Overall our clinics are performing well in controlling blood pressure in the >60-year-old population.

Age >= 60 with BP <150/90 * PCP Crosstabulation

			Clinic				Total
			DAN	IND	PG	SA	
Score	Fully met	Count	17	12	16	19	64
		% within Clinic	100.0%	100.0%	100.0%	100.0%	100.0%
Total		Count	17	12	16	19	64
		% within Clinic	100.0%	100.0%	100.0%	100.0%	100.0%

General Observations Applicable to all Clinics:

- A general good standard of care has been established.
- The use of coloured paper and other means of identifying different chart sections improves chart functionality.
- There is improved use of resources available, particularly in referrals to internists, nutritionists, and specialists.
- There is improved documentation as to why treatment goals are not met. For example, in cases where HbA1c is not met, there is documentation of non-compliance.
- NHI should consider performing an assessment on the quality of HbA1c offered to patient's in labs in the south. Many HbA1c are in the range of 4, which could either mean, quality of the test is not good, versus we are overmedicating the patients.
- For a counter-referral from a specialist another mechanism to receive it is to have an email address of the clinic attached to the original referral. The idea being that the specialist's services be advised of how to document a counter-referral from the clinic.
- It is advised that a training on the management of hypertension and its complications be provided to the physicians working for NHI. NHI could use this opportunity to update their guidelines regarding the management of the hypertensive patient
- Receiving specialists should have a channel to send counter referrals back to primary care physicians, to suggest changes to current management that can lead to improve patient outcomes.
 - Example from Hopkins files: A patient with abnormal ECG: it is noted that the patient is following a cardiologist, but there is no specific diagnosis noted from the cardiologist. The counter referral from the cardiologist is not found in the chart. The primary care provider shouldn't be satisfied with the patient's version about being followed by a cardiologist. Improved coordination with healthcare providers needs to be addressed.
- Over the years there has been an improvement in the overall documentation of files, however there is an issue that has been voiced repeatedly over the years regarding documentation. The physicians have noted that the requirement to document the visit in BHIS and NHI is very time consuming and redundant. As one physician pointed out, the time spent on double entry is time wasted. The time could be better spent tending to patients.

Observations per Clinic:

Dangriga and Satellites (Pomona, Hopkins) Clinical Audit:

Diabetes:

- Cases in which there has been increased HbA1c throughout the year have had the proper physical exercise and nutritional advice.
- The use of insulin and its use in difficult to manage patients has improved.
- There is appropriate referral to internist evaluation in challenging patients.
- Patient Note Comment: Patient with TC 190 mg/dL, LDL 87 mg/dL, HDL 46 mg/dL, Triglycerides 252 mg/dL and gets a prescription for gemfibrozil. BMI has been increasing in 2019. Obesity is not listed in the final diagnosis of the doctor after each encounter.
- Consider TSH testing in obese patients with a clinical syndrome suggestive of hypothyroidism (malaise, cold intolerance, difficulty losing weight, muscle pains, constipation)
- When experiencing cough with an ACE inhibitor, the GP should consider changing to an ARB instead of changing from one ace inhibitor to another

Hypertension:

- Stage 2 hypertension cases are getting good interdisciplinary care.
- Patient Note Comment: (SS#20089) Obese patient with a history of chest pains on 2016 for which a referral was given for patient to see cardiologist. No follow up on this referral on subsequent visits was noted and no ECG done since 2016.
- On analysis of ECG: We note that there is occasional copying of the ECG analysis provided by the electrocardiogram machine and that is written in the patient's notes. We should remember that these machines provide diagnosis that are based on measurements that the ECG machine takes. These machine diagnoses might not directly correspond to an actual clinical finding after a good history and physical examination.
- Patient Note Comment: (SSN 000233978) There is a history of CHF and diabetes. Patient is currently on treatment with amlodipine. According to guidelines ACE inhibitors or ARB have shown in different studies, to improve survival in patients with CHF, and in diabetics who have hypertension as a comorbidity.
 - CCB on the other hand, are second line in regard to hypertension treatment on a diabetic patient, once ACE inhibitors or ARB are available.
- Patient Note Comment: (000229253) 84-year-old female with a history of non-compliance and has chronic uncontrolled hypertension. She also has abnormal creatinine. No ECG has been done since 2014, no referral for internist assessment is noted. This type of patient is at risk due to age and comorbidities to suffer a stroke. The noncompliance issue should be discussed with family and reasons for noncompliance could be better documented.

Overall:

- Files are well organized.

- Overall the clinic's performance was very good. They have done a good job managing both hypertensive and diabetic cases.
- Patients with multiple comorbidities are receiving care that addresses their complex needs.
- Continuously document a patient's compliance or noncompliance and how it is affecting management.

Independence PC and Satellites Clinical Audit:

Diabetes:

- The use of insulin and its use in difficult to manage patients is good.
- There is appropriate referral to internist evaluation in challenging patients.
- There is good overall management.
-

Hypertension:

- Stage 2 hypertension cases are getting good interdisciplinary care.
- Patients with multiple comorbidities are receiving care that addresses their complex needs.
- Note ECG has not been functional for the last 4 months.
- Patient Note Comment: (SSN- 000247946) Obese female; fasting blood sugar (FBS) on 5th Feb 2019 144 mg/dL; fasting blood sugar on 8th March 2017 was 138 mg/dL; on 5th April 2018: fasting blood sugar was 187 mg/dL, urine test showed white blood cell too numerous to count (TNTC) and hemoglobin A1c 6.1. This patient is diagnosed as a hypertensive patient. Patient had multiple above normal FBSs and is obese. Although the HbA1c is normal, we should diagnose her as diabetic and get a second HbA1c repeated.
- Physicians should improve addressing increasing BMIs. Even though there's no nutritionist in the southern region, there is always something that can be done for the patient

Overall:

- Files are well organized.
- Overall the clinic's performance was very good. They have done a good job managing both hypertensive and diabetic cases.
- There is improved interdisciplinary collaboration noted.

Punta Gorda and Satellites Clinical Audit:

Diabetes:

- There is improved interdisciplinary collaboration noted.
- There is good overall management.
- Case Note (SSN 256733): Diabetic patient with weight loss and uncontrolled diabetes. Currently on metformin. Suggestions would be for her to be referred for internal medicine consultation for possible insulin therapy.
- Case Note (SSN 177076): Consider referring diabetic patient (age 38) with repeated admission/observation for hyperglycemia to social worker and internist to determine the underlying cause of repeated hyperglycemic episode.
- Case Note (SSN 418218): Diabetic patient: on Aug 3rd, 2018 an abdominal CT alerted to a tumoral mass in the pelvic cavity. On follow-up, March 3rd, 2019, patient is diagnosed with a high-risk pregnancy. Referral to ob-gyn was not noted and is suggested.

Hypertension:

- There is improved interdisciplinary collaboration noted.
- Stage 2 hypertension cases are getting good interdisciplinary care.
- Patients with multiple comorbidities are receiving care that addresses their complex needs.
- Patient Note Comment: (SSN - 110565) - Consider adding a secondary treatment to patient's treatment plan to help improve BP.
- Patient Note Comment: (SSN- 258081) - Consider switching patient from beta blocker to ACE or ARB to help improve patient's BP control, compliance and help provide renoprotection for patient as they are diabetic.
- Patient Note Comment: (SSN-6898): Diabetic, hypertensive patient with proteinuria. Currently on a CCB. Suggest switching or adding an ARB or ACE inhibitor for diabetic nephropathy protection.
- Patient Note Comment: (SSN – 413634). While BP is under control, the patients glucose value demonstrate that their DM is not controlled. Patient should be referred to an internist. Consider having patient see a social worker in the event the you believe the cause is noncompliance.

Overall:

- Files are well organized. Documentation of encounters is good. The use of colored paper for section identification is good.
- Overall the clinic's performance was very good. They have done a good job managing both hypertensive and diabetic cases.
- PG PCP referral consults for internal medicine go to Dangriga. Formal counter-referrals are not always available on patients return. We noted multiple cases with hyperglycemia that required urgent treatment. Timely referral for follow-up and possible insulin therapy initiation is suggested.
- Currently there is no nutritionist for formal nutritional consults.
- Observation: There appears to some form of 'cyclical compliance' whereby a patient's chronic condition appears to be under control for several visits followed by a period of being uncontrolled despite no change in treatment and back to being controlled. While this occurs throughout the

country, it appears that the prevalence of this is higher in Punta Gorda PCP and its satellite clinics. (Dr. Ajay Hotchandani)

San Antonio and Satellites Clinical Audit:

Diabetes:

- The use of insulin and its use in difficult to manage patients is good.
- There is appropriate referral to internist evaluation in challenging patients.
- SSN# 41631: A review of the Chronic Form shows the following trend of the microalbumin. A review of results should be done to determine if the data was properly imputed. If yes, then discuss the possibility of checking lab regarding accuracy.

Date	Microalbumin
14/10/16	80
16/8/17	10
14/2/18	80
18/12/18	10

Hypertension:

- There is improved interdisciplinary collaboration noted.
- Stage 2 hypertension cases are getting good interdisciplinary care.
- Patients with multiple comorbidities are receiving care that addresses their complex needs.

Overall:

- Files are well organized.
- Overall the clinic's performance was very good.
- They have done a good job managing both hypertensive and diabetic cases.
- Patients with multiple comorbidities are receiving care that addresses their complex needs.

Bibliography:

1. Jaramillo R, Ake C. *NHI Medical Audit South Side 2016 Report.*; 2016.
2. Jerums G, MacIsaac RJ. Treatment of microalbuminuria in patients with type 2 diabetes mellitus. *Treat Endocrinol.* 2002;1(3):163-173. <http://www.ncbi.nlm.nih.gov/pubmed/15799209>.
3. National Kidney Foundation. FREQUENTLY ASKED QUESTIONS ABOUT GFR ESTIMATES. https://www.kidney.org/sites/default/files/docs/12-10-4004_abe_faqs_aboutgfrrev1b_singleb.pdf. Accessed May 31, 2018.
4. Diabetic Retinopathy. American Optometric Association. <https://www.aoa.org/patients-and-public/eye-and-vision-problems/glossary-of-eye-and-vision-conditions/diabetic-retinopathy>. Published 2018.
5. Smoking and Diabetes | Overviews of Diseases/Conditions | Tips From Former Smokers | CDC. CDC. <https://www.cdc.gov/tobacco/campaign/tips/diseases/diabetes.html>. Published 2018.
6. PAHO. mhGAP: Drug Use and Drug Use Flowchart. https://www.paho.org/mhgap/en/drug_flowchart.html?reload#.
7. National Institute on Drug Abuse (NIDA). Nicotine. <https://www.drugabuse.gov/publications/brain-power/grades-6-9/legal-doesn%27t-mean-harmless-module-2/background>.
8. Verdecchia P, Angeli F, Mancina G, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension.* 2016;11(3):102-107. doi:10.1111/j.1365-2796.2011.02356.x
9. Jacobson TA, Ito MK, Maki KC, et al. National lipid association recommendations for patient-centered management of dyslipidemia: part 1--full report. *J Clin Lipidol.* 2015;9(2):129-169. doi:10.1016/j.jacl.2015.02.003
10. Nesto RW. LDL Cholesterol Lowering in Type 2 Diabetes: What Is the Optimum Approach? *Clin Diabetes.* 2008;26(1):8-13. doi:10.2337/diaclin.26.1.8
11. Jaramillo R. Proposal for Rationalization of Support Services Utilization (Final 2016) Done. 2016.
12. ASCVD Risk Estimator Plus - American College of Cardiology. <https://www.acc.org/tools-and-practice-support/mobile-resources/features/2013-prevention-guidelines-ascvd-risk-estimator>.
13. Long AN, Dagogo-Jack S. Comorbidities of diabetes and hypertension: mechanisms and approach to target organ protection. *J Clin Hypertens (Greenwich).* 2011;13(4):244-251. doi:10.1111/j.1751-7176.2011.00434.x
14. American Diabetes Association AD. Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2009;32 Suppl 1(Suppl 1):S62-7. doi:10.2337/dc09-S062
15. CDC. Defining Adult Overweight and Obesity | Overweight & Obesity | CDC. <https://www.cdc.gov/obesity/adult/defining.html>. Published 2018.
16. CDC. The Health Effects of Overweight and Obesity. CDC. <https://www.cdc.gov/healthyweight/effects/index.html>. Published 2018.
17. Onuigbo MAC, Agbasi N. Diabetic Nephropathy and CKD-Analysis of Individual Patient Serum Creatinine Trajectories: A Forgotten Diagnostic Methodology for Diabetic CKD Prognostication and Prediction. *J Clin Med.* 2015;4(7):1348-1368. doi:10.3390/jcm4071348
18. About Peripheral Artery Disease (PAD). http://www.heart.org/HEARTORG/Conditions/VascularHealth/PeripheralArteryDisease/About-Peripheral-Artery-Disease-PAD_UCM_301301_Article.jsp#.Wz5bptJKgU.
19. Clark CE, Taylor RS, Shore AC, Ukoumunne OC, Campbell JL. Association of a difference in systolic blood pressure between arms with vascular disease and mortality: a systematic review and meta-analysis. *Lancet (London, England).* 2012;379(9819):905-914. doi:10.1016/S0140-6736(11)61710-8
20. Liperoti R, Vetrano DL, Bernabei R, Onder G. Herbal Medications in Cardiovascular Medicine. *J Am Coll Cardiol.* 2017;69(9):1188-1199. doi:10.1016/j.jacc.2016.11.078

21. Rubenfire M. Herbal Medications in Cardiovascular Medicine - American College of Cardiology. American College of Cardiology. <https://www.acc.org/latest-in-cardiology/ten-points-to-remember/2017/03/02/12/27/herbal-medications-in-cardiovascular-medicine>. Published 2017.
22. *Belize Tobacco Control Report 2011.*; 2011. http://www.paho.org/hq/index.php?option=com_docman&task=doc_view&gid=16841&Itemid=270&lang=en.
23. Nhlbi. *Prevention, Detection, Evaluation, and Treatment of High Blood Pressure The Seventh Report of the Joint National Committee on Complete Report.*; 2004. <https://www.nhlbi.nih.gov/files/docs/guidelines/jnc7full.pdf>.